

## A. Creation Of 5' Nuclease Constructs

### 1. Modified DNAP $Taq$ Genes

5 The first step was to place a modified gene for the  $Taq$  DNA polymerase on a plasmid under control of an inducible promoter. The modified  $Taq$  polymerase gene was isolated as follows: The  $Taq$  DNA polymerase gene was amplified by polymerase chain reaction from genomic DNA from *Thermus aquaticus*, strain YT-1 (Lawyer *et al.*, *supra*), using as primers the oligonucleotides described in SEQ ID NOS:13-14. The resulting fragment of DNA has a recognition sequence for the restriction endonuclease EcoRI at the 5' end of the coding sequence and a BglII sequence at the 10 3' end. Cleavage with BglII leaves a 5' overhang or "sticky end" that is compatible with the end generated by BamHI. The PCR-amplified DNA was digested with EcoRI and BamHI. The 2512 bp fragment containing the coding region for the polymerase gene was gel purified and then ligated into a plasmid which contains an inducible promoter.

15 In one embodiment of the invention, the pTTQ18 vector, which contains the hybrid *trp-lac* (*tac*) promoter, was used [M.J.R. Stark, *Gene* 5:255 (1987)] and shown in Fig. 14. The *tac* promoter is under the control of the *E. coli lac* repressor. Repression allows the synthesis of the gene product to be suppressed until the desired level of bacterial growth has been achieved, at which point repression is removed by 20 addition of a specific inducer, isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG). Such a system allows the expression of foreign proteins that may slow or prevent growth of transformants.

25 Bacterial promoters, such as *tac*, may not be adequately suppressed when they are present on a multiple copy plasmid. If a highly toxic protein is placed under control of such a promoter, the small amount of expression leaking through can be harmful to the bacteria. In another embodiment of the invention, another option for repressing synthesis of a cloned gene product was used. The non-bacterial promoter, from bacteriophage T7, found in the plasmid vector series pET-3 was used to express the cloned mutant  $Taq$  polymerase genes [Fig. 15; Studier and Moffatt, *J. Mol. Biol.*

189:113 (1986)]. This promoter initiates transcription only by T7 RNA polymerase. In a suitable strain, such as BL21(DE3)pLYS, the gene for this RNA polymerase is carried on the bacterial genome under control of the *lac* operator. This arrangement has the advantage that expression of the multiple copy gene (on the plasmid) is completely dependent on the expression of T7 RNA polymerase, which is easily suppressed because it is present in a single copy.

For ligation into the pTTQ18 vector (Fig. 14), the PCR product DNA containing the *Taq* polymerase coding region (*mutTaq*, clone 4B, SEQ ID NO:21) was digested with *EcoRI* and *BglIII* and this fragment was ligated under standard "sticky end" conditions [Sambrook *et al. Molecular Cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, pp. 1.63-1.69 (1989)] into the *EcoRI* and *BamHI* sites of the plasmid vector pTTQ18. Expression of this construct yields a translational fusion product in which the first two residues of the native protein (Met-Arg) are replaced by three from the vector (Met-Asn-Ser), but the remainder of the natural protein would not change. The construct was transformed into the JM109 strain of *E. coli* and the transformants were plated under incompletely repressing conditions that do not permit growth of bacteria expressing the native protein. These plating conditions allow the isolation of genes containing pre-existing mutations, such as those that result from the infidelity of *Taq* polymerase during the amplification process.

Using this amplification/selection protocol, we isolated a clone (depicted in Fig. 4B) containing a mutated *Taq* polymerase gene (*mutTaq*, clone 4B). The mutant was first detected by its phenotype, in which temperature-stable 5' nuclease activity in a crude cell extract was normal, but polymerization activity was almost absent (approximately less than 1% of wild type *Taq* polymerase activity).

DNA sequence analysis of the recombinant gene showed that it had changes in the polymerase domain resulting in two amino acid substitutions: an A to G change at nucleotide position 1394 causes a Glu to Gly change at amino acid position 465 (numbered according to the natural nucleic and amino acid sequences, SEQ ID NOS:1 and 4) and another A to G change at nucleotide position 2260 causes a Gln to Arg

change at amino acid position 754. Because the Gln to Gly mutation is at a nonconserved position and because the Glu to Arg mutation alters an amino acid that is conserved in virtually all of the known Type A polymerases, this latter mutation is most likely the one responsible for curtailing the synthesis activity of this protein. The nucleotide sequence for the Fig. 4B construct is given in SEQ ID NO:21. The enzyme encoded by this sequence is referred to as Cleavase® A/G.

Subsequent derivatives of DNAP*Taq* constructs were made from the mut*Taq* gene, thus, they all bear these amino acid substitutions in addition to their other alterations, unless these particular regions were deleted. These mutated sites are indicated by black boxes at these locations in the diagrams in Fig. 4. In Figure 4, the designation "3' Exo" is used to indicate the location of the 3' exonuclease activity associated with Type A polymerases which is not present in DNAP*Taq*. All constructs except the genes shown in Figures 4E, F and G were made in the pTTQ18 vector.

The cloning vector used for the genes in Figs. 4E and F was from the commercially available pET-3 series, described above. Though this vector series has only a BamHI site for cloning downstream of the T7 promoter, the series contains variants that allow cloning into any of the three reading frames. For cloning of the PCR product described above, the variant called pET-3c was used (Fig 15). The vector was digested with BamHI, dephosphorylated with calf intestinal phosphatase, and the sticky ends were filled in using the Klenow fragment of DNAPEc1 and dNTPs. The gene for the mutant *Taq* DNAP shown in Fig. 4B (mut*Taq*, clone 4B) was released from pTTQ18 by digestion with EcoRI and SalI, and the "sticky ends" were filled in as was done with the vector. The fragment was ligated to the vector under standard blunt-end conditions (Sambrook *et al.*, *Molecular Cloning, supra*), the construct was transformed into the BL21(DE3)pLYS strain of *E. coli*, and isolates were screened to identify those that were ligated with the gene in the proper orientation relative to the promoter. This construction yields another translational fusion product, in which the first two amino acids of DNAP*Taq* (Met-Arg) are